

Please, note that for reasons of shortness only the allergenic content is stated on the product label.

V.2 (< 051029>)



BIOMAY AG

Vienna Competence Center
Lazarettgasse 19 · Top1
A-1090 Vienna · AUSTRIA

Tel: +43/1/79 66 296-100

Fax: +43/1/79 66 296-111

e-mail: info@biomay.com

www.biomay.com

Hev b 6.02-MBP

(Hevein domain of Allergen 6 from *Hevea brasiliensis* (Latex),

Fusion to Maltose binding protein)

For research purpose only.

PRODUCT DESCRIPTION:

Access:EMBL M36986/Swissprot: P02877

Mw = 47536 Dalton

Mol. Ext. Coeff.: 78400; 1mg/ml $A_{278} = 1.65$

pI = 5.02

Lot#: 01


Amount: 1 mg

Quality: Purity $\geq 97\%$

Reacts with IgE from Hev b 6.02 reactive human serum

General Information

BIOMAY Hev b 6.02-MBP is a recombinant fusion protein consisting of Maltose binding protein (MBP) and the Hevein domain of Hev b 6 (43 AA) fused to the C-terminal end of MBP.



The construct contains a factor Xa cleaving site that allows the isolation of the Hev b 11 fragment. It was produced by heterologous expression in *E. coli*, purified by conventional biochemical methods, and lyophilized from 5 mM NH_3HCO_3 (volatile).

Reconstitution:

The material should be reconstituted with distilled water (or equivalent) or dilute buffers. Do not use salt concentrations exceeding 20 mM to dissolve the lyophilized material. Salt may be added after

dissolution. Gentle agitation during dissolution is essential, afterwards incubate for 30 min to allow a complete reconstitution of the protein. If reconstituted to 1 mg/ml the product will be soluble to at least 98%.

Storage:

When stored at -20°C the quality of the material will be maintained for several years. However, for short periods (max. 3 weeks) the lyophilized product may be kept at room temperature. After reconstitution store at -20°C . Avoid repeated freezing/thawing.

Quality control:

Purity has been determined on SDS-PAGE gels stained with Coomassie Brilliant Blue R-250.

Hev b 6.02-MBP tested positive in an IgE-Immunoblot using a standardized pool of human Hev b 6.02-reactive sera.

* The mol.ext.coeff. was calculated from the DNA-derived protein sequence as described by Gill, S.C. and by Hippel, P.H. (1989) Analytical Biochemistry **182**, 319-326.