

Please, note that for reasons of shortness only the allergenic content is stated on the product label.

V.2 (<051029>)



BIOMAY AG

Vienna Competence Center
Lazarettgasse 19 · Top1
A-1090 Vienna · AUSTRIA

Tel: +43/1/79 66 296-100
Fax: +43/1/79 66 296-111
e-mail: info@biomay.com
www.biomay.com

Hev b 9.1 –His

(Allergen 9 Isoform 1, Enolase from *Hevea brasiliensis*, His-tagged)

For research purpose only.

PRODUCT DESCRIPTION:

Access: EMBL: AJ132580/Swissprot: Q9LEJO

M_w = 47,830 Dalton

Mol. Ext. Coeff.: 43,654; 1mg/ml A₂₇₈=0.91*

pI = 5.6

Lot#: 01

Amount: 1 mg

Quality: Purity better than 98%

General Information:

BIOMAY Hev b 9.1- His is expressed in *E. coli* as a His-tagged protein. The protein was purified by Ni²⁺ affinity chromatography and ion exchange chromatography. The product was lyophilized in 5 mM NH₄HCO₃ (volatile) containing 1% sucrose.

Reconstitution:

The material can be reconstituted with water or diluted buffers. If reconstituted with water or buffers (2mM β-Mercaptoethanol) to 2 mg/ml sucrose concentration will be app. 0.6%.

If reconstituted with water or buffers (2mM β-Mercaptoethanol) to 2 mg/ml, the product is soluble to app. 99%. Thorough physical suspension of the protein is essential. Alternatively the product can be dissolved in 6M Urea, 1mM β-Mercaptoethanol. The urea solution can be dialyzed against a suitable buffer (20 mM

Tris/HCl pH 8.0 1mM β-Mercaptoethanol) without precipitation of protein.

Storage:

The lyophilized product can be kept at room temperature for at least 2 weeks. However, we recommend the product to be stored at -20°C. Under these conditions the quality of the material will be maintained for several years. The stability at 4°C should at least be 6 months. Reconstituted protein can be stored at -20°C.

Quality control:

By SDS-PAGE and staining with Coomassie-blue R250. (Immunological properties were controlled by SDS-PAGE\Western-blotting with Hev b 9- specific human IgE)

* The mol.ext.coeff. was calculated from the DNA-derived protein sequence as described by Gill, S.C. and by Hippel, P.H. (1989), Analytical Biochemistry **182**, 319-326.