

Phl p 6 (Phl p 6.0101)

(*Phleum pratense*, timothy grass pollen allergen 6)

The leader sequence of the protein is replaced by a methionine

For research purpose only.

PRODUCT DESCRIPTION:

Access: EMBL Z27082 /Swissprot P 43215

Mw = 11,790 Dalton

Mol. Ext. Coeff.: 5,960; 1 mg/ml $A_{280} = 0.506^*$

pI = 5,56

Lot#: 03

Amount: 1 mg / 250 µg

Quality: Purity ≥ 99%

Endotoxin content: 0,0002 EU/µg

Reacts with IgE from Phl p 6-reactive human serum.

BIOMAY AG

Vienna Competence Center
Lazarettgasse 19 Top 1
A-1090 Wien

Tel: +43 1 7966296-0
Fax: +43 1 7966296-111
e-mail: info@biomay.com
www.biomay.com



General information:

BIOMAY Phl p 6 is a recombinant protein with IgE-binding capacity. It was produced by heterologous expression in *E. coli*, purified by conventional biochemical methods, and lyophilized from sodium phosphate buffer (pH 7.4)

Reconstitution:

The material should be reconstituted with distilled water (or equivalent) or dilute buffers. Do not use salt concentrations exceeding 20 mM to dissolve the lyophilized material. Salt may be added after dissolution. Gentle agitation during dissolution is essential, afterwards incubate for 30 min to allow a complete reconstitution of the protein. If reconstituted by this procedure the product will be soluble to a concentration of at least 1 mg/mL. At this protein concentration the concentration of sodium phosphate will be 2 mM.

Storage:

When stored at -20°C the quality of the material will be maintained for several years. However, for short periods (max. 3 weeks) the lyophilized product may be kept at room temperature. After reconstitution store at -20°C. Avoid repeated freezing/thawing.

Quality control:

Purity has been determined on SDS-PAGE gels stained with Coomassie Brilliant Blue R-250. Endotoxin content was determined by using a Limulus Amebocyte Lysate (LAL) assay. The above stated lot tested positive in an IgE-Immunoblot with human Phl p 6-reactive serum.

* The mol.ext.coeff. was calculated from the DNA-derived protein sequence as described by Gill, S.C. and by Hippel, P.H. (1989), Analytical Biochemistry **182**, 319-326.